

**Coherence in Category Theory and the
Church-Rosser Property**

by

C. Barry Jay

Coherence in Category Theory

LFCS Report Series

ECS-LFCS-91-181

LFCS

October 1991

Department of Computer Science
University of Edinburgh
The King's Buildings
Edinburgh EH9 3JZ

Copyright © 1991, LFCS

Coherence in Category Theory and the Church-Rosser Property ^{*}

C. Barry Jay [†]

LFCS, Department of Computer Science
University of Edinburgh
The King's Buildings, Mayfield Road
Edinburgh, UK, EH9 3JZ
e-mail: cbj@dcs.ed.ac.uk

October 22, 1991

Abstract

Szabo's derivation systems on sequent calculi with exchange and product are not Church-Rosser. Thus his coherence results for categories having a symmetric product (either monoidal or cartesian) are false.

1 Introduction

Gentzen's sequent calculi [9] have been applied extensively in category theory, e.g. [2, 3, 4, 6, 7, 8]. Sequents correspond to morphisms of a category, and the rules of the calculus correspond to categorical structures (e.g. having an associative tensor product). Cut-elimination was then used to put bounds on the complexity of these structures, e.g. to produce exhaustive lists (perhaps with duplications) of the canonical natural transformations between given functors. For symmetric, monoidal closed categories it was shown in [12] how to decide in principle whether two such transformations are equal, while an effective, linear-time decision procedure was given in [1].

Derivation systems (reduction rules) can be used to eliminate some duplicates in the list of cut-free proofs (e.g. [8]). However, in Algebra of Proofs [11] and its forerunner [10], Szabo claims to have produced derivation systems in which all

^{*}To appear in Notre Dame Journal of Formal Logic

[†]Research supported by The Royal Society of Edinburgh/BP, and NSERC operating grant OGPIN 016.

duplicates have been eliminated, so that every proof has a unique normal form. In fact, none of the systems there which include a symmetry for the product are Church-Rosser (confluent).

The two major applications of this work were complete coherence theorems for symmetric, monoidal closed categories (mentioned above) and for cartesian closed categories. The former claim leads to self-contradictory assertions about the size of homsets in free categories. The latter problem is solved in [5] using similar methods, but avoids adopting symmetry as a primitive by exploiting the universal property of the cartesian product.

I would like to thank Professor S. Mac Lane for raising this issue and exhorting me to resolve it.

2 The Counter-example

Here is Szabo's sequent calculus for symmetric, monoidal categories [11]

$$\frac{\Gamma \rightarrow \gamma \quad \Delta \gamma \Lambda \rightarrow \phi}{\Delta \Gamma \Lambda \rightarrow \phi} (Cut) \qquad \frac{\Gamma \Delta \rightarrow \phi}{\Gamma \Lambda \Delta \rightarrow \phi} (LI) \qquad \frac{\Gamma \alpha \beta \Delta \rightarrow \phi}{\Gamma \beta \alpha \Delta \rightarrow \phi} (Ex)$$

$$\frac{\Gamma \rightarrow \alpha \quad \Delta \rightarrow \beta}{\Gamma \Delta \rightarrow \alpha \otimes \beta} (R\otimes) \qquad \frac{\Gamma \alpha \beta \Delta \rightarrow \phi}{\Gamma \alpha \otimes \beta \Delta \rightarrow \phi} (L\otimes)$$

which are there labelled (R1), (R2),(R4),(R8) and (R9), respectively. Here is a proof in the calculus:

$$\frac{\frac{\frac{\frac{\frac{\frac{\alpha \rightarrow \alpha \quad \beta \rightarrow \beta}{\alpha \beta \rightarrow \alpha \otimes \beta} (R\otimes)}{\alpha \beta \gamma \rightarrow (\alpha \otimes \beta) \otimes \gamma (= \phi)} (R\otimes)}{\alpha \gamma \beta \rightarrow \phi} (Ex)}{\alpha \beta \gamma \rightarrow \phi} (Ex)}{\beta \alpha \gamma \rightarrow \phi} (L\otimes)}{\beta (\alpha \otimes \gamma) \rightarrow \phi} (Ex)}{(\alpha \otimes \gamma) \beta \rightarrow \phi} (L\otimes)}{\alpha \otimes \gamma \otimes \beta \rightarrow \phi} (L\otimes)} \frac{\gamma \rightarrow \gamma}{\alpha \beta \gamma \rightarrow (\alpha \otimes \beta) \otimes \gamma (= \phi)} (R\otimes) (Ex)$$

Two derivations (reductions) applicable to this proof are:

(D.24.1) "For any permutation π of the integers $1, \dots, n$, and any successions (σ) and (τ) of instances of R4,

$$\frac{\Gamma \rightarrow \bar{\Phi}}{\Delta \rightarrow \bar{\Phi}} (\sigma) \qquad \geq \qquad \frac{\Gamma \rightarrow \bar{\Phi}}{\Delta \rightarrow \bar{\Phi}} (\tau)$$

provided that $\Gamma = \alpha_1 \dots \alpha_n$, $\Delta = \beta_1 \dots \beta_n$, $\beta_i = \alpha_{\pi(i)}$ for $1 \leq i \leq n$, and (τ) is the unique string of interchanges which first moves $\alpha_{\pi(1)}$ to β_1 , then $\alpha_{\pi(2)}$ to β_2 , etc. if π is the identity permutation, the right-hand side denotes [the null proof]."

(D.27.3)

$$\frac{\frac{\frac{\Gamma\alpha\beta\gamma\Delta\rightarrow\Phi}{\Gamma\alpha\gamma\beta\Delta\rightarrow\Phi}}{\Gamma\gamma\alpha\beta\Delta\rightarrow\Phi}}{\Gamma\gamma(\alpha\otimes\beta)\Delta\rightarrow\Phi} \geq \frac{\frac{\Gamma\alpha\beta\gamma\Delta\rightarrow\Phi}{\Gamma(\alpha\otimes\beta)\gamma\Delta\rightarrow\Phi}}{\Gamma\gamma(\alpha\otimes\beta)\Delta\rightarrow\Phi}$$

The first two applications of (Ex) in this proof can be deleted by (D.24.1) to yield an irreducible proof. However, (D.27.3) can also be applied, followed by (D.24.1) to obtain a distinct irreducible proof. Thus the system is not Church-Rosser. Categorically, the two irreducible proofs correspond to the morphisms

$$ca(c\otimes 1) = a^{-1}(1\otimes c)a : (\alpha\otimes\gamma)\otimes\beta \rightarrow (\alpha\otimes\beta)\otimes\gamma$$

whose proof of equality requires an identity morphism to be expanded to c^2 .

The loss of confluence leads to incorrect combinatorial calculations of the size of homsets. Let \mathcal{V} be a free symmetric monoidal closed category on a set of objects. For objects A and B of \mathcal{V} let $\|A, B\|$ denote the cardinality of $\mathcal{V}(A, B)$. Further, let $A^{(1)} = [A, I]$ denote the dual of A and $A^{(n)}$ its n th dual. Claim (2) of Corollary 8.6.13 is that if A is not isomorphic to I we have

$$\|A^{(2n)}, A^{(2m)}\| = \binom{n+m-1}{n} \|A, A\|$$

This formula is self-contradictory since it implies that

$$\|A^{(4)}, A^{(4)}\| = \binom{3}{2} \|A, A\| = 3 \|A, A\|$$

while applying it twice shows that

$$\|A^{(4)}, A^{(4)}\| = \|A^{(2)}, A^{(2)}\| = \|A, A\|$$

Similar calculations show that claims (3),(4) and (5) are also false.

References

- [1] C.B. Jay, "The structure of free closed categories", J. Pure Appl. Alg., 66 (1990) 271-285.
- [2] G.M. Kelly and S. Mac Lane, "Coherence in closed categories", J. Pure Appl. Alg. 1 no. 1, (1971), 97-140; erratum, ibid. no. 2, 219.
- [3] J. Lambek, "Deductive systems and categories I. Syntactic calculus and residuated categories", Math. Systems Theory 2 (1968) 287-318.

- [4] J. Lambek, "Deductive systems and categories II. Standard constructions and closed categories", in: *Lecture Notes in Mathematics* 86 (Springer-Verlag, 1969) 76–122.
- [5] J. Lambek and P. Scott, *Introduction to higher order categorical logic*, Cambridge studies in advanced mathematics 7, Cam. Univ. Press (1986).
- [6] S. Mac Lane, "Why commutative diagrams coincide with equivalent proofs", *Contemporary Mathematics* 13 (1982) 387–401.
- [7] G.E. Minc, "A cut elimination theorem for relevant logics", *J. Soviet Math.* Vol 6 no 4 (1976), 422–428, translated from *Zap. Nauch Sem. Leningrad. Otdel. Mat. Inst. Steklov (LOMI)* 32 (1972), 90–97.
- [8] G.E. Minc, "Closed categories and the theory of proofs", *J. Soviet Math.* 15 No. 1 (1981) 45–62, translated from *Zap. Nauch Sem. Leningrad. Otdel. Mat. Inst. Steklov (LOMI)* 68 (1977), 83–114.
- [9] M.E. Szabo (Ed.), *The collected papers of Gerhard Gentzen*, Studies in Logic and the Foundations of Mathematics, North-Holland, Amsterdam (1969).
- [10] M.E. Szabo, "A categorical equivalence of proofs", *Notre Dame J. Formal Logic* XV No. 2 (1974) 177–91.
- [11] M.E. Szabo, *Algebra of Proofs*, North-Holland, Amsterdam (1978).
- [12] R. Voreadou, "Coherence and non-commutative diagrams in closed categories", *Mem. Amer. Math. Soc.* No. 182 (1977).

**Copyright © 1991, Laboratory for Foundations of Computer Science,
University of Edinburgh. All rights reserved.**

**Reproduction of all or part of this work
is permitted for educational or research use
on condition that this copyright notice is
included in any copy.**